



# E-Cigarette Use Among Youth and Young Adults

A Report of the Surgeon General

## Fact Sheet

This Surgeon General's report comprehensively reviews the public health issue of e-cigarettes and their impact on U.S. youth and young adults. Studies highlighted in the report cover young adolescents (11-14 years of age); adolescents (15-17 years of age); and/or young adults (18-25 years of age). Scientific evidence contained in this report supports the following facts:

**E-cigarettes are a rapidly emerging and diversified product class. These devices typically deliver nicotine, flavorings, and other additives to users via an inhaled aerosol. These devices are referred to by a variety of names, including "e-cigs," "e-hookahs," "mods," "vape pens," "vapes," and "tank systems."**

- E-cigarettes are battery-powered devices that heat a liquid into an aerosol that the user inhales.
- The liquid usually has nicotine, which comes from tobacco; flavoring; and other additives.
- E-cigarette products can also be used as a delivery system for marijuana and other illicit drugs.

**E-cigarettes are now the most commonly used tobacco product among youth, surpassing conventional cigarettes in 2014. E-cigarette use is strongly associated with the use of other tobacco products among youth and young adults, including cigarettes and other burned tobacco products.**

- In 2015, more than 3 million youth in middle and high school, including about 1 of every 6 high school students, used e-cigarettes in the past month. More than a quarter of youth in middle and high school have tried e-cigarettes.
- Among high school students, e-cigarette use is higher among males, whites, and Hispanics than among females and African-Americans.
- There is a strong association between the use of e-cigarettes, cigarettes, and the use of other burned tobacco products by young people. In 2015, for example, nearly 6 of 10 high school cigarette smokers also used e-cigarettes.
- Research has found that youth who use a tobacco product, such as e-cigarettes, are more likely to go on to use other tobacco products like cigarettes.

**E-cigarette use among youth and young adults has become a public health concern. In 2014, current use of e-cigarettes by young adults 18-24 years of age surpassed that of adults 25 years of age and older.**

- Among young adults 18-24 years of age, e-cigarette use more than doubled from 2013 to 2014. As of 2014, more than one-third of young adults had tried e-cigarettes.
- The most recent data available show that the prevalence of past 30-day use of e-cigarettes was 13.6% among young adults (2014) and 16.0% among high school students (2015).
- The most recent data available show that the prevalence of past 30-day use of e-cigarettes is similar among middle school students (5.3%) and adults 25 years of age and older (5.7%).
- Among young adults, e-cigarette use is higher among males, whites and Hispanics, and those with less education.

**The use of products containing nicotine poses dangers to youth, pregnant women, and fetuses. The use of products containing nicotine in any form among youth, including in e-cigarettes, is unsafe.**

- Many e-cigarettes contain nicotine, which is highly addictive.
- The brain is the last organ in the human body to develop fully. Brain development continues until the early to mid-20s. Nicotine exposure during periods of significant brain development, such as adolescence, can disrupt the growth of brain circuits that control attention, learning, and susceptibility to addiction.
- The effects of nicotine exposure during youth and young adulthood can be long-lasting and can include lower impulse control and mood disorders.
- The nicotine in e-cigarettes and other tobacco products can prime young brains for addiction to other drugs, such as cocaine and methamphetamine.